




The Effect of Dynamic Variable Resistance (DVR) on Muscular Strenght

The first concept of Variable Resistance Exercise Machines introduced by Dr. Gideon Ariel



Code	adi-pub-01108
Title	The Effect of Dynamic Variable Resistance (DVR) on Muscular Strenght
Subtitle	The first concept of Variable Resistance Exercise Machines introduced by Dr. Gideon Ariel
Name	Computerized Biomechanical Analysis Inc.
Author	Gideon Ariel
Published on	Monday, March 1, 1976
Subject	ACES; Exercise Machine; Favorite; Journal; Legal; Media; Science; Sports; Studies; Track and Field
URL	https://arielweb.com/articles/show/adi-pub-01108
Date	2013-01-16 15:40:45
Label	Approved
Privacy	Public

Universal's Dynamic Variable Resistance Doubles Strength Gains Over Olympic Bar Training

An independent study by Computerized Biomechanical Analysis Incorporated, commissioned by Universal Gym Equipment, compared the average strength gains from two different resistive weight training systems - Universal's Dynamic Variable Resistance and the Olympic Barbell.

The study involved twenty male University students with weight training experience of at least two years. The experiment was conducted over a twenty week period, with the subjects divided into two groups. Group 1 trained on conventional Olympic barbell equipment, while Group 2 trained on dynamic variable resistance (D.V.R.) exercise equipment.

The results showed that both groups increased their mean strength level during the twenty week period. However, the group using the Olympic barbell increased their strength level from 249.5 pounds to 285.5 pounds, demonstrating a mean change of 36 pounds. On the other hand, the dynamic variable resistance (D.V.R.) exercise group increased their mean strength in the bench press exercise from 252.5 pounds to 327.0 pounds, demonstrating a mean change of 74.5 lbs.

The study concluded that training the muscle in a dynamic fashion alone, such as with Olympic bar lifting, does not cause maximum efficiency of muscular training. When varying the resistance according to biomechanical principles, muscular gains are greater, both practically and statistically.

This PDF summary has been auto-generated from the original publication by arielweb-ai-bot v1.2.2023.0926 on 2023-09-28 03:40:04 without human intervention. In case of errors or omissions please contact our aibot directly at ai@macrosport.com.

Copyright Disclaimer


The content and materials provided in this document are protected by copyright laws. All rights are reserved by Ariel Dynamics Inc. Users are prohibited from copying, reproducing, distributing, or modifying any part of this content without prior written permission from Ariel Dynamics Inc. Unauthorized use or reproduction of any materials may result in legal action.

Disclaimer of Liability


While every effort has been made to ensure the accuracy of the information presented on this website/document, Ariel Dynamics Inc. makes no warranties or representations regarding the completeness, accuracy, or suitability of the information. The content is provided "as is" and without warranty of any kind, either expressed or implied. Ariel Dynamics Inc. shall not be liable for any errors or omissions in the content or for any actions taken in reliance thereon. Ariel Dynamics

Below find a reprint of the 16 relevant pages of the article "The Effect of Dynamic Variable Resistance (DVR) on Muscular Strenght" in "Computerized Biomechanical Analysis Inc.":

HERE'S PROOF!
UNIVERSAL'S DYNAMIC
VARIABLE RESISTANCE



DOUBLES YOUR
STRENGTH GAINS
OVER OLYMPIC BAR
TRAINING.



© Universal Gym Equipment, 1977

The following independent study by
Computerized Biomechanical Analysis
Incorporated was commissioned by
Universal Gym Equipment to compare
the average strength gains from two
different resistive weight training
systems — Universal's Dynamic Variable
Resistance and the Olympic Barbell.

Universal GYM EQUIPMENT
1328 N. Sierra Vista • Fresno, Ca. 93703 • Phone: (209) 251-4251
A subsidiary of First Northwest Industries of America, Inc.

The Effect of Dynamic Variable Resistance (D.V.R.) on Muscular Strength

By
Gideon B. Ariel, Ph.D.

March, 1976

**Computerized
biomechanical
analysis Incorporated**

Science serving industry, sports and human performance

316 College Street • Amherst, Mass. 01002 • Phone: 413-256-0486

THE EFFECT OF DYNAMIC VARIABLE RESISTANCE (D.V.R.) EXERCISE ON MUSCULAR STRENGTH

Introduction

The relationship between resistance exercises and muscle strength has been known for centuries. In ancient Greece, Milo, the Greek wrestler, used progressive resistance exercises to improve his strength. His original method consisted of lifting a calf each day until its full growth, and this technique provides the first example of progressive resistance exercises.

The 434 voluntary muscles in man constitute 40 to 60 percent of his total body weight. These muscles are responsible for human motion, which is the most fundamental function of the musculoskeletal system.

Muscular strength may be defined as the force a muscle group can exert against a resistance in a maximal effort. In 1948, Delorme (4) adopted the name "progressive resistance exercise" for his method of developing muscular strength through the utilization of counterbalancing and weight of the extremity with a cable and pulley arrangement and, thus, gave load-assisting exercise to muscle groups which did not perform antigravity motions. McQueen (6) distinguished between exercise regimens for producing muscle hypertrophy and those for producing muscle power. He concluded that the number of repetitions for each set of exercise determines the different characteristics of the exercise. Based on evidence presented in these early studies, hundreds of investigations have been published relative to muscular development including isotonic exercises, isometric

exercise, eccentric contraction techniques, Oxford technique, double and triple progressive super set system, and many others. Each system has been supported and refuted by numerous investigations. Berger (3) concluded that 6-7 repetitions three times a week is best for developing dynamic strength. Other research conducted by Steinhaus (9) emphasized the need to increase the intensity—not the amount of work—in order to develop maximum strength.

The most recent research pertaining to exercise was conducted by Thomas B. Pipes and Jack H. Wilmore (7) in their article contrasting isokinetic with isotonic strength training in adult men. According to their findings with isokinetic contractions of both in low speed and high speed contractions, the results demonstrated a clear superiority of the isokinetic training procedure over the isotonic procedures. In 1972, Ariel (8) introduced the Dynamic Variable Resistance exercise principles which resulted in variable resistance exercise equipment. For the first time biomechanical principles were employed in the design of exercise equipment, and rather than force a man to fit the machine, the machine was designed for the man.

In sport and athletics, most movements are ballistic in nature. This implies that they are preprogrammed as a unit in the central mechanisms of the brain and, once initiated, cannot be influenced by sensory and/or environmental information. This necessitates exact precision in the timing and coordination of both the system of muscle contraction, as well as, in the segmental sequence of muscular activity involved with complex tasks.

In order to accomplish ballistic movement, it is necessary to utilize isotonic exercise routines. It is impossible to duplicate the neuromuscular system utilizing isokinetic exercises which by their nature control the speed of the movement. It was found that a characteristic pattern of motion is present during intentional movement of body segments against resistance. This pattern consists of reciprocally organized activity between the agonist and antagonist. These reciprocal activities occur in consistent temporal relationships with motion parameters, such as velocity, acceleration, and forces. Hellebrandt and Houtz (5) shed some light on the mechanism of muscle training in an experimental demonstration of the overload principle. They found that mere repetition of contractions which place little stress on the neuromuscular system had little effect on the functional capacity of the skeletal muscles; however, they found that the amount of work done per unit of time is the critical variable upon which extension of the limits of performance depends. The speed with which functional capacity increases suggests that central nervous system, as well as the contractile tissue, is an important contributing component of training.

Since the human body is a system of linked segments, forces cause rotation of these segments about their anatomical axes. Both muscle and gravitational forces are important in producing these turning effects which are fundamental in body movements in all sports and daily living. Pushing, pulling, lifting, kicking, running, walking, and all human activities are results of rotational motion of the links which are made of bones. Since force has been considered the most important component of

athletic performance, many exercise equipment manufacturers have developed various types of devices employing isometrics and isokinetics. These isometric and isokinetic devices inhibit the natural movement patterns of acceleration and deceleration. However, when considered as a separate entity, force is only one factor influencing successful athletic performance.

The three factors underlying all athletic performance are:

1. Force

2. Displacement (direction of movement)

3. Duration of movement

In all motor skills, muscular forces interact to move the body parts through the activity. The displacement of the body parts and their speed of motion are important in the coordination of the activity and are also directly related to the forces produced. However, it is only because of the control provided by the brain that the muscular forces follow any particular displacement pattern, and without these brain center controls, there would be no skilled athletic performances. In all athletic events, the intricate timing of the varying forces is a critical factor in successful performances, and therefore, training an isolated muscle group slowly or at a constant speed may result in poorer athletic performances.

THE DYNAMIC VARIABLE RESISTANCE (D.V.R.) CONCEPT:

In conventional resistive exercises, loads are moved through a range of motion. The muscular force and the load are not constant because of the modifying effects of the lever system throughout the range of motion. In an

exercise such as the bench press, there is a point where the resistance is maximum and below or above this point the resistance is less. This fact illustrates the important phenomenon that throughout an exercise stroke, the muscle is working at its maximum potential during a very small range of motion.

To facilitate maximum muscular involvement it is necessary to vary the resistance. In some exercises, this resistance should vary by as much as 100 percent in order to maintain the moment of force at its maximum. The resistance should be varied according to biomechanical data obtained under dynamic conditions.

The purpose of the present study was to compare the new dynamic variable resistance (D.V.R.) exercise concept to a conventional resistance training method.

METHOD

Twenty male University students between the ages of 19 and 23 were used in the present study. Their height averaged 181.5 cm. with a mean weight of 91.4 kg. The experiment was conducted during a twenty week period.

All subjects were athletes with weight training experience of at least two years. For a period of four weeks prior to the beginning of the test, all the subjects lifted weights five days each week and were tested on the seventh day of the week for maximum lifts in the bench and military presses, the curl, and the squat. A standard warm-up procedure was performed and each test was a maximal lift. The experiment was conducted during a subsequent twenty week period.

The twenty subjects were divided into two equal groups. Those subjects in Group 1 were assigned to train on conventional Olympic barbell equipment, while those in Group 2 were assigned training procedures on dynamic variable resistance (D.V.R.) exercise equipment.*

Subjects trained five times per week for approximately two hours each training session, according to a program designed to work the major muscle groups of the body utilizing a progressive over-load principle. The program consisted of bench press, military press, squat, and curl exercise. Each exercise was performed in sets of four with an increased load following a pyramidal increase. Each set consisted of a decreasing number of repetitions from 8 to 3. Weights were increased as rapidly as possible to maintain the training at maximum effort. The training programs for both groups were identical with the only difference being the exercise equipment. Constant supervision of workouts was maintained at all times. Testing was conducted every six days until the conclusion of the study. Both experimental groups were tested for maximum dynamic strength on the Olympic bench press set. This exercise was selected due to the similarity of the procedures. The bench press exercise was conducted as the subject reclined on his back flat against the bench. The weight was handed to the subject, and then lowered by the subject to his chest. Immediately, the barbell was raised to a straight arm position directly above the chest.

*Apparatus used in this experiment was provided by Universal Gym Equipment. Centurion - Dynamic Variable Resistance Machines, Fresno, California.

Muscular force testing was conducted during the four weeks prior to the start of the experimental period and at six day intervals thereafter until the conclusion of the study. Maximum dynamic muscular force measurements were determined by 1-RM in the bench press. Techniques used in performing the press were those prescribed by the Amateur Athletic Union rules for weight-lifting competition.

The reader should notice that maximum strength testing was conducted on the Olympic bench press set even though only Group 1 trained with the Olympic barbells. Group 2 trained only with the dynamic variable resistance apparatus. This factor introduced a bias against those subjects training with the dynamic variable resistance since these individuals were not repeatedly exposed, via their training regimen, to the Olympic bars.

RESULTS

As voluntary muscular force gains are a function of training over a period of time, neither the "t-test" nor a one-way analysis of variance for "before" and "after" conditions were deemed totally adequate analytic techniques. The relationship between both the gains in muscular force and the effect of time on the two experimental groups suggested a comparison of the regression lines to investigate the influences of the two different types of exercise equipment upon the development of muscular strength. Therefore, analysis of variance and slope analysis were utilized in assessing the data.

Table 1 presents the raw data for the bench press exercise. Table 2 illustrates the means, standard deviations and the variances. Figure 1 illustrates the mean changes in the muscular strength in the bench press exercise during the twenty week experimental period. Table 3 illustrates the regression analysis results between the two experimental groups.

It was found that both groups increased their mean strength level during the twenty week period. The group using the Olympic barbell increased their strength level from 249.5 pounds to 285.5 pounds demonstrating a mean change of 36 pounds. However, this strength increase was not found to be statistically significant, although such a strength gain may represent a practical significance. The dynamic variable resistance (D.V.R.) exercise group increased their mean strength in the bench press exercise from 252.5 pounds to 327.0 pounds demonstrating a mean change of 74.5 lbs which was statistically significance, as well as, practical significance.

Table 4 revealed the analysis of variance between the beginning and the last period.

Regression analysis yielded a significant difference between the slope of the two regression lines. The regression coefficient for the Olympic barbell group was 1.31, illustrating an average gain of 1.31 pounds per week while the dynamic variable resistance (D.V.R.) group produced regression coefficient of 3.84 demonstrating an average gain of 3.84 pounds per week. Comparison of the slopes of the regression lines yielded a significant F-ratio demonstrating the statistically significant difference between the two training methods in favor of the dynamic variable resistance exercise (D.V.R.) method.

Subject	1st Week	2nd Week	3rd Week	4th Week	5th Week	6th Week	7th Week	8th Week	9th Week	10th Week	11th Week	12th Week	13th Week	14th Week	15th Week	16th Week	17th Week	18th Week	19th Week	20th Week
A	175	175	180	180	185	185	180	190	195	200	205	210	210	220	220	225	230	230	235	240
B	175	175	180	180	185	185	180	190	195	200	205	210	210	220	220	225	230	230	235	240
C	205	205	210	210	210	210	215	215	215	215	215	220	220	220	220	215	225	230	230	235
D	205	210	215	220	225	225	225	235	245	245	250	250	255	260	260	265	265	270	275	280
E	215	215	225	225	225	230	230	230	235	240	245	245	240	245	250	250	250	250	255	255
F	215	215	225	235	245	255	255	260	265	270	270	270	275	275	275	285	285	290	295	295
G	220	220	220	225	225	225	230	230	230	235	235	240	245	245	250	250	255	260	260	260
H	220	230	230	240	245	250	255	260	265	270	275	280	280	285	290	290	295	300	305	310
I	220	225	225	225	230	230	235	240	245	250	255	260	265	270	275	280	280	285	290	290
J	230	230	235	240	245	250	255	265	275	285	290	295	300	310	315	315	320	325	330	340
K	230	235	240	240	240	245	250	250	250	255	260	265	265	270	270	270	270	270	275	275
L	235	235	240	245	250	255	260	265	270	270	280	285	285	290	290	300	305	305	310	320
M	265	265	270	270	275	280	280	280	285	275	285	285	290	295	295	300	295	300	295	295
N	275	280	300	315	315	320	320	325	325	330	335	335	340	340	340	345	350	350	350	355
O	300	300	300	315	315	320	320	320	320	320	325	320	325	325	315	320	320	320	325	325
P	300	300	310	310	315	325	325	325	330	330	335	340	340	345	350	350	350	350	350	355
Q	325	325	330	330	340	345	345	350	350	350	350	350	345	350	355	355	360	360	360	360
R	330	335	340	340	340	350	350	350	350	355	360	365	370	370	375	375	375	375	375	380
S	340	340	345	345	345	340	340	345	350	350	350	350	350	355	355	355	355	360	360	360
T	340	340	345	355	355	360	360	365	365	370	375	375	380	380	380	380	375	380	385	385

Olympic Bar Group in black numbers
*The two groups, each contained 10 men, were matched for similar or equal starting strengths.

Universal DVR (Dynamic Variable Resistance) Group in red numbers
starting strengths.

	1st Week	2nd Week	3rd Week	4th Week	5th Week	6th Week	7th Week	8th Week	9th Week	10th Week	11th Week	12th Week	13th Week	14th Week	15th Week	16th Week	17th Week	18th Week	19th Week	20th Week
Mean	249.50	250.00	254.00	256.50	258.50	261.50	263.00	265.00	267.00	267.00	270.00	271.00	273.50	275.00	275.00	278.50	281.50	283.00	284.50	285.50
Variance	3058.05	3038.88	3021.11	3122.50	3233.61	3139.16	3076.88	3144.44	3190.00	3106.84	3105.55	3251.11	3328.89	3073.33	3329.16	3035.83	3043.33	2943.33	3090.00	3105.55
St. Dev.	55.29	55.12	54.96	55.87	56.86	56.02	55.43	56.07	56.48	55.88	55.72	56.86	57.72	55.46	57.81	55.85	55.77	54.16	55.12	55.13

Olympic Bar Group in black numbers
Universal DVR (Dynamic Variable Resistance) Group in red numbers

DISCUSSION

The results of the present experiment demonstrate that training the muscle in a dynamic fashion alone, such as with Olympic bar lifting, does not cause maximum efficacy of muscular training. When varying the resistance according to biomechanical principles, muscular gains are greater, both practically and statistically.

When a subject uses a resistance device such as the Olympic barbell, there are two kinds of forces applied to this system. The internal forces produced by the muscular system and the external forces produced by the resistance device, in this case, the weight or the bar. When considering any human force system, muscles, bones, and joints, as well as externally applied resistance, must be

considered. The actual forces produced by individual muscles cannot be predicted easily because of the indeterminate influence of a number of physiological and mechanical factors. These include length-tension and force-velocity relationships, as well as, the location of the muscle attachments with respect to the joint and the dynamic effect of the movement. It is to examine the muscular involvement in the exercise to determine the moments of force produced by all the muscles at the particular joint. It is well known in resistance exercises that there exists a "sticking point" during which the apparent resistance is at its maximum. However, the absolute muscular force is relatively constant and varies slightly depending on the force-length relationship.

DISCUSSION
The results of the present experiment demonstrate that training the muscle in a dynamic fashion alone, such as with Olympic bar lifting, does not cause maximum efficiency of muscular training. When varying the resistance according to biomechanical principles, muscular gains are greater, both practically and statistically. When a subject uses a resistance device such as the Olympic barbell, there are two kinds of forces applied to this system. The internal forces produced by the muscular system and the external forces produced by the resistance device, in this case, the weight or the bar. When considering any human force system, muscles, bones, and joints, as well as externally applied resistance, must be

considered. The actual forces produced by individual muscles cannot be predicted easily because of the indeterminate influence of a number of physiological and mechanical factors. These include length-tension and force-velocity relationships, as well as, the location of the muscle attachments with respect to the joint and the dynamic effect of the movement. One way to determine the muscular involvement in the exercise is to examine the moments of force produced by all the muscles at the particular joint. It is well known in resistance exercises that there exists a "sticking point" during which the apparent resistance is at its maximum. However, the absolute muscular force is relatively constant and varies slightly depending on its force length relationship.

TABLE 3
COMPARISON OF REGRESSION LINES BETWEEN
BARBELL AND THE VARIABLE EXPERIMENTAL GROUPS

	d.f.	Reg. Coef.	Standard Error	F-ratio (between regression lines)
Barbell Group	9	1.87	.61	
D.V.R. Group	9	3.84	1.15	18.65** (1.17)

**F-ratio significant at the .01 level of confidence.

TABLE 4
ANALYSIS OF VARIANCE FOR PROGRESSION
IN STRENGTH LEVELS

OLYMPIC BAR GROUP				F-ratio
Week 1	1	10	249.5	
Week 20	2	10	285.5	
	1	6460	6460	2.344718267
	10	49745	2763.611111	
	19	56225	52.5705967	
			Mean difference = 36.0 lbs.	

DYNAMIC VARIABLE RESISTANCE GROUP				F-ratio
Week 1	1	10	252.5	
Week 20	2	10	327	
	1	27751.25	27751.25	10.47821071**
	10	47672.5	2648.472222	
	19	75423.75	51.46330948	
			Mean difference = 74.5 lbs.	

** Significant to the .01 level of confidence.

UNIVERSALS' DYNAMIC VARIABLE RESISTANCE
VS. CONVENTIONAL OLYMPIC BAR
IN THE BENCH-PRESS EXERCISE


The graph plots Strength Levels (lbs. x 10) on the Y-axis (ranging from 20 to 32) against Weeks on the X-axis (ranging from 1 to 20). Two lines represent the data: a red line for the D.V.R. Group and a black line for the Barbell Group. Both groups show an upward trend, with the D.V.R. Group consistently higher than the Barbell Group throughout the 20-week period.

Weeks	D.V.R. Group (lbs. x 10)	Barbell Group (lbs. x 10)
1	25.5	25.0
2	26.0	25.0
3	26.5	25.0
4	27.0	25.0
5	27.5	25.0
6	28.0	25.0
7	28.5	25.0
8	29.0	25.0
9	29.5	25.0
10	30.0	25.0
11	30.5	25.0
12	31.0	25.0
13	31.5	25.0
14	32.0	25.0
15	32.5	25.0
16	33.0	25.0
17	33.5	25.0
18	34.0	25.0
19	34.5	25.0
20	35.0	25.0

FIGURE 1

In conventional resistance exercise such as on the bench press, loads are moved through a range of motion. The mass remains constant throughout the motion but the muscular forces and the dynamic forces are not constant because of the modifying effects of the lever system throughout the range of motion. For all practical purposes, the absolute muscular force is the same throughout the exercise since the only difference is the force arm

UNIVERSAL'S DYNAMIC VARIABLE RESISTANCE VS. CONVENTIONAL OLYMPIC BAR IN THE BENCH-PRESS EXERCISE



In conventional resistance exercise such as on the bench press, loads are moved through a range of motion. The mass remains constant throughout the motion but the muscular forces and the dynamic forces are not constant because of the modifying effects of the lever system throughout the range of motion. For all practical purposes, the absolute muscular force is the same throughout the exercise since the only difference is the force arm

on which the muscle pulls. When the force arm changes due to the angular changes of the limb, the muscle can lift a variable load. This explains why when performing an exercise such as the bench press, there is a point where the resistance is maximum and below or above this point the resistance is less. This fact illustrates the important phenomenon that throughout an exercise stroke using conventional Olympic barbells, the muscle is working at its maximum potential during a very small range of motion.

Another factor to consider in muscular training is the dynamic characteristics of the motion. In conventional barbell lifting, there is an initial burst of muscular activity as the agonist muscle contracts and the antagonist muscle relaxes thus causing acceleration of the limb. This is followed by an intervening quiet period during which there is no muscular firing activity and which is followed by deceleration of the limb as the antagonist contracts. Near the end of the movement, the antagonist muscle has to stop the motion. With a conventional barbell, the stopping motion starts too soon causing a diminished training effect. The dynamic variable resistance exercise equipment assigns different resistances throughout the range of motion in order to accommodate the biomechanical changes occurring during the exercise and, at the same time, adjusts for the ballistic characteristics of the movement. With the dynamic variable resistance apparatus, the agonist muscle can fire for a longer period of time. This type of ballistic training is shown by the results of the present study to be more efficient for dynamic muscular training.

BIBLIOGRAPHY

1. Ariel, G. Computerized Biomechanical Analysis of Human Performance. Mechanics and Sport. New York: ASME Press, 1973.
2. Ariel, G. Computerized Biomechanical Analysis of Track and Field Athletics. Track and Field Quarterly Review 72: 99-103, 1973.
3. Berger, R. Effect of Varied Weight Training Programs on Strength. Research Quarterly 33: 168-181, 1962.
4. Delorme, T. Restoration of Muscle Power with Resistance Exercise. J. Bone and Joint Surg. 27: 645-651, 1945.
5. Hellebrandt, F. and Houtz, S. Mechanism of Muscle Training in Man: Experimental Demonstration of Overload Principle. Physical Therapy Review 36: 371-376, 1956.
6. McQuenn, I. Recent Advances in the Technique of Progressive Resistance Exercise. Brit. Med. J. 2: 328-338, 1954.
7. Pipes, T., and Wilmore, J., Isokinetic vs isotonic strength Training in Adult Men. Medicine and Science in Sports. 7: 262-274, 1976.
8. Ariel, G. Computerized Biomechanical Analysis of the Variable Resistance Exercise Machine. Technical Report, UNI-7, 1974.
9. Steinhaus, H. Strength of Morpurgo to Muller - A Half Century of Research. J. Assoc. Physical And Mental Rehab. 9: 147-150, 1955.